
H B 1323-M6 9F636m 1949

Florida, State Board of Health. Bureau of Vital Statistics.

Maternal Mortality



NI M OFOLLOEN D

NLM 05011354 4

NATIONAL LIBRARY OF MEDICINE

ARMY MEDICAL LIBRARY

FOUNDED 1836



WASHINGTON, D.C.

the tables who people at their report of makening materially an address of Charles I interested property alter faction of themselves to maken the address of the control of

to being there were his trademandation occurs to the real time of the party of the


Harting delicate and the second of the secon

Table I place a segretar of the property of the property of the party of the party

Table 2 sizes the Christian Confidence and the Confidence of the C

The parties without the property of the parties of

The to be a local to the control of the second of the seco



State Board of Health BUREAU OF VITAL STATISTICS Jacksonville, Florida

MATERNAL MORTALITY IN FLORIDA

August 1, 19 49

The tables and graphs of this report on maternal mortality are designed to acquaint all interested persons with Florida's standing in regard to maternal deaths. Maternal deaths include all deaths as a result of conditions arising from pregnancy, childbirth, and the puerperlum.

In 1948, there were 115 resident maternal deaths in Florida and the death rate was 1.9 per thousand live births. This is the lowest maternal death rate on record in this state and it represents a decrease of 70% since 1940. The colored maternal death rate of 4.2 was almost 4 times as high as the white rate of 1.1 per thousand live births.

Florida's maternal death rate is still higher than the National average. Data issued by the National Office of Vital Statistics for the year 1947 (the latest year for which final data for all states has been released) showed Florida as being in 44th place when compared with other states. Only four states (Georgia, Alabama, Missiasippi, and South Carolina) had a higher maternal mortality rate, and all of the above are neighthoring states.

Table A gives a comparison of maternal mortality rates (per 1,000 live births), by color, for Florida and the United States from 1933 to 1948. The rate for the United States in 1948 is a provisional figure, since final data has not yet been made awalilable, by the National Office of Vital Statistics.

Graph 1 shows the trend in maternal mortality rates in Florida and the United States from 1933 to 1948, while Graph 2 shows the trend in maternal mortality rates in Florida, by color, for the same period. These graphs are based upon the data contained in Table A.

Table B gives the live births distributed according to the age of the mother, maternal deaths according to age of death, and maternal mortality rates by age groups. The largest number of births for both white and colored was in the 20-24 age group which also had the lowest maternal mortality rate. The highest maternal mortality rate for both white and colored was in the 35-39 age group.

The resident maternal mortality rate (per 1,000 live births) for Florida in 1944-1948 was 2.6. Charlotte County had the lowest rate with no maternal deaths while Liberty County had the highest rate of 10.7. Table C gives the number of maternal deaths for each year from 1944 to 1948, the number of live births for the combined period, and the maternal mortality for each county for 1944-1948 arranged in order of magnitude.

The maternal mortality rate in Table C is given for the combined period of 1944-1948 because rates based upon a small number of events must be interpreted with caution. Even with the combined period, the reader must keep an open mind in interpreting data in this table.

Table D gives the maternal deaths by cause, by year, by color, together with the corresponding maternal mortality rates. The combined maternal deaths and maternal mortality rates are also shown for the entire period. Puerperal Toxemias (List No. 148) was the leading cause of maternal deaths for 1944-1948 while infection during Childbirth and the Puerperlum (List No. 147) was the second highest cause.



WATERNAL MORTALITY IN FLORIDA

Will of States

es and graphs of this report on maternal margingley are dealined to nequent to parkens alth FlorigaC c. standing it regard to maternal dealine. Maternal dealine as a result of conditions origing from prognammy, childbirth,

there noted 225 rould one autornal deaths in Floride and the death rate on the this test lies alorest rate on the this conference of FD & since 1940. The colored natural rote rate of the since 1940. The colored natural rote rate of the thousand of the pirthes.

place a consertion of maternal mortality rates that 1,000 time of ends, place in the ottends, place and the United States for 1933 to 1948. The rese for the United States and participant flyers, place that date has not get peen date washingle at Ortice of Vital testicales.

anous the trent in neturnal northilty rates in Signide, and the United anough to the United in maternal northing rates by the to the tend in maternal northing rates by select for the same period. These graphs are breed upon the data con-

gives the live sirths distributed according to the age of the mothers the according to age according to both abits and colored ear in the 20-24 age group abigh legest material martality rate. The bighost deternal martality rate. For

The sections of the same and the coloner than the section of the s

ernal mortality rate in Table 2 is given for the consined period of 1944rates naced upon a small number of events must be interpreted with acutions a consined period, the reader numb keep as open wind in interpreting date in

qives the motornol deaths by caves, by paor, by color, together with the sontenest motorial motorial serves. The sontened meternal deaths and natural metals are also shown for the soil of pariods 'unique of Towenlan illier no. 1962 in a cause of maternal deaths for 1982-1982 white integrine during Chitablett sportum illet her last mater accord highest mouses.

TABLE A. RESIDENT MATERNAL MORTALITY RATES (Per 1,000 Live Births)
BY COLOR, FLORIDA AND THE UNITED STATES, 1933 - 1948

YEAR		FLORIDA		UNI	TED STAT	ES
TEAN	TOTAL	WHITE	COLORED	TOTAL	WHITE	COLORED
1948	1.9	1.1	4. 2	1.2*		
19 47	2.2	1.3	5.0	1.3	1.1	3.3
1946	2.9	1.9	5.9	1.6	1.3	3.6
19 45	2.9	1.8	6.0	2.1	1.7	4.3
1944	3.3	2.3	6.0	2.3	1.9	5.1
1943	3.5	2.7	6.1	2.5	2.1	5.1
1942	4.0	2.5	8.0	2.6	2.2	5.4
1941	6.1	4.5	10.0	3.2	2.7	6.8
1940	6.4	5.0	9.8	3.8	3.2	7.7
1939	6.4	5.5	8.5	4.0	3.5	7.6
19 38	7.6	5.5	12.6	4.4	3.8	8.5
1937	6.7	5 • 2	10.1	4.9	4.4	8.6
1936	7.9	6.2	12.0	5-7	5.1	9.7
19 35	8.6	7.3	11.7	5.8	5.3	9.5
1934	8.3	6.9	11.6	5.9	5.5	9.0
1933	11.1	8.9	15.9	6.2	5.7	9.7

^{*}Provisional data. Not available by color.

TABLE 8. RESIDENT BIRTHS, MATERNAL DEATHS, AND MATERNAL MORTALITY RATES (Per 1,000 Live Births) BY AGE GROUPS, BY COLOR, FLORIDA, 1948

AGE OF		BIRTH	5	MAT	ERN AL DE	ATHS	MATE	RN AL MORT	FALITY
MOTHER	TOTAL	WHITE	COLORED	TOTAL	WHITE	COLORED	TOTAL	WHITE	COLORED
10-14	208	43	165	0	0	0	-	-	-
15-19	9,571	5,684	3,887	18	6	12	1.9	1. 1	3.1
20-24	19,729	14,673	5,056	21	7	14	1.1	0.5	2.8
25-29	15,239	11,763	3,476	28	10	18	1.8	0.9	5.2
30-34	8,880	6,829	2,051	20	. 9	11	2.3	1.3	5.4
35-39	4,698	3,410	1,288	24	12	12	5.1	3.5	9.3
40-44	1,127	811	316	4	2	2	3.5	2.5	6.3
45-49	77	39	38	0	0	0	-	-	-
50 and	3	2	1	0	0	0	-	-	-
Unknown	153	82	71	0	0	0	-	-	-
Al I Ages	59,685	43, 336	16,349	115	46	69	1.9	1.1	4.2

RESIDENT MATERNAL MORTALITY RATES (FAT 1,000 11me SIRTAD) BY COLOR, FLORIDA AND THE UNITED STATES, 1933 - 194

	-					
	TATE OFFI			AUTEURS		
	37109					
			5.0			
		2.3			2.1	
			0.0		0.0	
E =#						
I.e.				Tub		
				6.5		
0,1			+511011			
Est .	146					
			0.0	215		
2.40			12.6			
			2.02	8+8		
8.0				2+3		
				Est	7.9	
	9+9					
949		2+2			(+8	
					Lell	


crolos es siccifico for estab lenslateor

STORNT BIRTHS, MATERNAL DEATHS, AND MATERNAL MORTALITY RATES

	out the			146 26V	AS INI			5 70
	ATRON JAM							
		MATOT				COLORED	HTHI	-
								1.200
15.00								gr.
								1. 13
			1 172					1 65
917							A. EDV.II	FEE
								20
								88
								VE
								11
								E
								8.88
	Let		69					- 11
								8 16


GRAPH 1

RESIDENT MATERNAL MORTALITY (Per 1,000 Live Births), FLORIDA AND THE UNITED STATES, 1933-1948




RATE PER 1,000 LIVE BIRTHS

RESIDENT MATERNAL MORTALITY (Per 1,000 Live Births),



CHARACT

ACCEPTANT MATERIAL MORTALITY (For 1,000 List elected), FLORIDA



RESIDENT MATERIAL MORTALITY FORF 1,000 LIVE DIRECTOR





TABLE C. RESIDENT MATERNAL DEATHS (For Each Year), LIVE BIRTHS, AND MATERNAL MORTALITY RATES (Per 1,000 Live Births), BY COUNTY, FLORIDA, 1944-1948


COUNTIES		MAT	ERNAL DEA	THS		LIVE BIRTHS	MATERNAL MORTALITY
DOUNT I E S	1944	1945	19 46	1947	19 48	1944 - 1948	1944 - 194
FLORIDA	160	142	156	135	115	272, 258	2.6
1. Charlotte	0	0	0	0	0	345	-
2. Brevard	1	. 0	0	0	1	2,383	0.8
3. Calhoun	0	0	1	0	0	1,178	0.8
4. Taylor	C	0	0	1	0	1,178	0.8
5. Franklin	1	0	0	0	0	853	1.2
6. Santa Rosa	0	1	0	2	0	2,330	1.3
7. Orange	3	3	2	4	3	10,583	1.4
8. Wakulla	0	1 0	0	0	1	638	1.6
9. Martin	0	0	0	1	0	626	1.6
C. Indian River	1	0	1	0	0	1,209	1.7
1. Manutee	0	2	0	2	1	3,004	1.7
2. Monroe	2	. 0	0	1	1	2.305	1.7
3. Hillsboro	15	4	10	9	10	26,659	1.8
4. St. Lucie	2	0	0	2	0	2, 163	1.8
5. Sarasota	3	0	1	1 0	0	2, 173	1.8
6. Dade	19	13	19	9	12	38,518	1.9
7. Hardee	0	1	0	0	1	1,034	1.9
8. Seminole	1	1	3	1	0	2,992	2.0
9. Holmes	1 0	1	1	1	1	1,936	2.1
C. Walton	1	1	1	0	1	1,875	2.1
1- Clay	1	0	. 0	1	1	1,387	2.2
2. Highlands	1 0	2	1	1	. 0	1,809	2. 2
3. Polk	6	! 8	3	5	1	12,527	2.2
4. Sumter	1	1 0	1	1	0	1,264	2.4
5. Pinellas	4	E	7	5	: 1	10,523	2.4
6. St. Johns	1	2	2	1	. 0	2,498	2.4
7. Palm Beach	6	4	6	3	6	10,347	2.4
3. Duval	13	19	20	16	15	33,718	2.5
9. Washington	1	1 0	1	2	. 0	1,621	2.5
D. Lafayette	1	0	0	0	, 0	40 2	2.5
l. Escambla	: 9	12	8	: 5	3	14,407	2.6
2. Bay	. 2	4	į 4	2	2	5,219	2.7
3. Oksechobee	1	lo lo	0	. 0	C	372	2.7
4. Alachua	1 3	3	3	2	: 5	5,710	2.8
5. Flagler	1	0	0	! 0	, 0	349	2.9
6. Volusla	1 4	5	2	4	3	6,239	2.9
7. Okaloosa	1	1	2	2	1 3 1	3,078	2.9
3. Nassau	2	0	2	. 1	0	1,653	3.0
o osceola	0	0	1	1 2	1	987	3.0
. Broward	1 3	1	5	1 7	6	6,997	3.1
1. Lee	. 2	1	3	. 0	2	2,470	3.2
2. Citrus	. 0	1	1	0	1 0	600	3.3
3. Lake	. 4	1 3	3	. 2	0	3,416	3.5
4. Bradford	1	1	0	2	2	1,634	3.7
5. Baker	1	0	2	. 0	. 0	815	3.7
6. Leon	3	5	14	5	0	4,936	3.8
. Madison	3	1	2	2	0	2,037	3.9
. Jackson	1 5	5	3	4	2	4, 787	4.0
. Hernando	1	. 0	1	1	0	751	4.0
. Pasco	; 0	1	3	2	1	1,746	4.0
1. Columbia	5	3	:	0	2	2,461	4.1 4.1
2. DeSoto	2	1	0	1	C	9 78	
3. Putnam	2	: 2	2	1	3	2,216	4.5
4. Glades	0	: 0	0	0	1	221	4.5
5. Suwannee	4	3	0	2	1	2,194	4.7
6. Gulf	0	0	0	2	3	1,065	4.7
7. Dixle	1	0	0	2	0 1	1,216	4.9
B. Hamilton	1 0	4	1	3.	4	3,934	5.3
9. Harlon	1 3	3	6	5	1	741	5.4
. Hendry	1	0	1	2	0	1,227	5.7
1. Levy 2. Union	3 2	1 2	0	3	0 1	677	5.9
3. Gadsden	2	4	6	3	11	3,756	6.9
4. Glichrist	1	1	1	3	Ö	410	7.3
5. Jefferson	2	2	2	3	2	1,404	7.8 8.8
6. Collier	. 0	1	2	0	1	456	10.7

HARRIST SALE STATE OF THE PARTY

TABLE D	D. RESIDENT MATERNAL DEATHS AND	0 0	ATH	RATES	(Per	000 %	Live	irths	BY CAL	USE, B	100 Y	OR, FL	ORIDA,	35 44-19 48
2014	=			MATERN	AL DE	S H L		5 Year		oc	ATE	10		5 Year
LI 36 MO.	n n D 4		1944	1945	1946	1947	1948	Total	1944	1946	19 46	1947	1943	Rate
34		⊢¥∪	165 68 74	142 66 75	156	135	11 00 00 00 00 00 00 00 00 00 00 00 00 0	333 375	50 00 00 00 00 00 00 00 00 00 00 00 00 0	2.00	20.00	2000	1.2	7.6
140	Abortion with mention of infection	⊢ ≥ ∪	125	10	14 7 7	100	20 ± 02	25	0.30	0.20		0.26	0.10	0.20
141	Abortion without mention of infection	⊢ ≥ ∪	sc 는 다	F 10 (4)	r 3 E	. 60 W W	E 4 8	13	0.10	0.14	0.13	0.13	0.05	0.11 0.06 0.19
142	Ectopic gestation	ب≆ں	ott	amo	111	400	871	200	0.08	, 8 8	0.20	0.07	0.013	0.13
143	Hemorrhage of pregnancy (death before delivery)	ل≥ ن	E0 H	m-1 0	000	000	нон	Ø 167 3	000	0.09	0.05	0 0	0.00	0000
144	Toxemias of pregnancy (death before delivery)	⊢ ≥ ∪	19	E ~ 8	11 9	10	12	63 63		0.31	000		0.20	
145	Other diseases and accidents of pregnancy (death before delivery)	⊢≩U	ran	000	240	505	tot	19 5	0.00	0.04	0.06	0.08	0.07	0.02
146	Hemorrhage of childbirth and the puerperlum	⊢≩U	23	19	1178	10	125	109	0 0 0 1	w 40	0 0 0			
147	Infection during childbirth and the puerporium	F B U	33	26	704	700	18	123	0.67	0.57	0.44	0.33	0.30	1.00
148	Puerperal toxemiss fexcluding death before delivery)	⊢ ≱ ∪	30	26	29	20	23	128	0.61	0.53	0.53	0.33	0.39	0.30
149	Other accidents and specified conditions of childbirth	F 3.0	10 10	# 10 #	17	26 10 16	700	3.5.0	0.28	0.29	0.31	0.43	0.23	0.31
150	Other and unspecified conditions of childsirth and the puerperium	- ≥ 0	a to	O 10 3	α) a≯ a≯	E 10 80	20 50 50	2 2 2 t	0.18	0.18	0.15	0.22	0.13	0.17







PRESSBOARD
PAMPHLET BINDER

Manufactured by
GAYLORD BROS. Inc.
Syracuse, N. Y.
Stockton, Calif.

JUL 2 7 1850

HB 1323.M6 qF636m 1949

01650230R

NLM 05011354 4

NATIONAL LIBRARY OF MEDICINE